

Caring for your child at home after a **COLONOSCOPY**

Dear Parents,

Your child has just had a colonoscopy. Colonoscopy is a procedure that involves passing a scope (flexible tube) through the rectum, and into the large intestine. It is to observe the lining (wall) of those structures. A biopsy from the lining of the intestine may also be done. This is normally not painful because the mucosal lining of the intestinal tract does not have nerve endings. This following information will help ensure your child has a safe recovery after leaving the hospital.

Care after the procedure

- A small amount of rectal bleeding (no more than a teaspoon) can be expected after the procedure.
- Your child may complain of abdominal fullness or discomfort. This is due to the air that was injected into the
 gastro-intestinal tract during the procedure. Walking or lying down on the right side is some of the ways to get
 relief from this discomfort.

Pain management

- Give acetaminophen (Tylenol® or Tempra®) every 4 hours if needed, but no more than 5 doses total in 24 hours. Nextdosemay begiven only at _____hours.
- Other medications might be prescribed. It is very important to follow the exact dosage and frequency prescribed.
- Refer to other instructed pain management information.

When will life go back to normal?

- Activity: Quiet activities today with close supervision. Normal activities tomorrow.
- Bathing: May have a bath or shower tomorrow.
- **Diet:** Give water, apple juice, Popsicle, flat ginger ale or Jell-O for one to two hours. If tolerated, proceed to a soft diet such as soup, yogurt, ice cream and/or cereal, and then advance to a normal diet.
- School/Daycare: May go back to school/daycaretomorrow.

When to present to the emergency room?

- Pain that does not improve even after giving your child the acetaminophen.
- A fever of 38.5° C (101° F) or higher, within 24 hours after the procedure.
- Rectal bleeding persists or more than one teaspoon of blood.
- Abdominal pain or bloating that gets worse.
- Any change or increase in the original symptoms.

Follow-up

- Your gastroenterologist will call you with the biopsy results in 3-4 weeks.
- Further follow-up will be arranged depending on the results

Disclaimer: Information provided by this pamphlet is for educational purposes. It is not intended to replace the advice or instruction of a professional healthcare practitioner, or to substitute for medical care. Contact a qualified healthcare practitioner if you have any questions concerning the care of your child.